

MATRIX FACTORIZATIONS AND DOUBLE LINE IN $\mathfrak{sl}_n$ QUANTUM LINK INVARIANT

YASUYOSHI YONEZAWA

Dedicated to Professor Akihiro Tsuchiya on the occasion of his retirement

ABSTRACT. This article gives matrix factorizations for the trivalent diagrams and double line appearing in $\mathfrak{sl}_n$ quantum link invariant. These matrix factorizations reconstruct Khovanov-Rozansky homology. And we show that the Euler characteristic of the matrix factorization for a double loop equals the quantum dimension of the representation $\wedge^2 V$ of $U_q(\mathfrak{sl}_n)$ in Section 3.2.

CONTENTS

1. Introduction	1
2. Category of matrix factorization	3
2.1. Matrix factorization	3
2.2. $\mathbb{Z}$ -graded matrix factorization	5
2.3. Koszul matrix factorization	6
3. Matrix factorization for trivalent diagrams and double line	9
3.1. Khovanov Rozansky homology	9
3.2. Definition of matrix factorization for trivalent diagrams and double line	11
References	17

1. INTRODUCTION

L. Kauffman introduced a graphical link invariant which is the normalized Jones polynomial [4][5]. It is well-known that the polynomial is derived from the fundamental representation of the quantum group $U_q(\mathfrak{sl}_2)$. Further, G. Kuperberg constructed a graphical link invariant associated with the fundamental representation of the quantum group $U_q(\mathfrak{sl}_3)$ [10]. H. Murakami, T. Ohtsuki and S. Yamada introduced a graphical regular link invariant for the fundamental representation of the quantum group $U_q(\mathfrak{sl}_n)$ [11]. In general, we can also obtain a graphical link invariant for a given quantum group $U_q(\mathfrak{g})$ and the fundamental representation. These invariants are collectively called $\mathfrak{g}$ quantum link invariants.

M. Khovanov constructed a categorification of $\mathfrak{sl}_2$ quantum link invariant [6]. A categorification generally means the replacement of a set with a category by corresponding an element to an object. The morphism of the category is properly chosen to carry theory well-done. For a categorification, there is an inverse operation called a decategorification which is the replacement of a category with a set. The decategorification of equivalent objects in the category is a same element in the set.

The Khovanov's theory is a beautiful example of the categorification; this is the replacement of Jones polynomial $J(L)$, which is a map from the set of links to " $\mathbb{Z}[q, q^{-1}]$ ", by a map C_K from the set of links to "the homotopy category of the bounded complex of graded $\mathbb{Z}$ -modules". The bounded complex $C_K(L)$ and the $\mathbb{Z} \oplus \mathbb{Z}$ -graded homology groups $KH^{i,j}(L)$ associated with $C_K(L)$ also become link invariants under the Reidemeister

moves. The decategorification is a $\mathbb{Z}$ -graded Euler characteristic χ_{KH} with the normalized Jones polynomial $J(L)$;

$$\chi_{KH} \left(\bigoplus_{i,j \in \mathbb{Z}} KH^{i,j}(L) \right) := \sum_{i,j \in \mathbb{Z}} (-1)^i \dim_{\mathbb{Q}}(KH^{i,j}(L) \otimes \mathbb{Q}) q^j = J(L).$$

Recently, M. Khovanov and L. Rozansky introduced a categorification of $\mathfrak{sl}_n$ quantum link invariant using the homotopy category of the bounded complex of matrix factorizations [7][8]. Since a resolution of a link diagram consists of a combination of the two local diagrams  and  (See FIGURE 2), a matrix factorization for the resolution is defined by a tensor product of some matrix factorizations for the two local diagrams. Then, the categorification of $\mathfrak{sl}_n$ quantum link invariant is constructed as the map $\mathcal{C}_{KR}$ from the set of links to the homotopy category of the bounded complex of matrix factorizations for some resolutions.

The bounded complex $\mathcal{C}_{KR}$ and the $\mathbb{Z} \oplus \mathbb{Z} \oplus \mathbb{Z}_2$ -graded homology groups $KRH^{i,j,k}(L)$ associated with $\mathcal{C}_{KR}(L)$ also become link invariants under the Reidemeister moves. The Euler characteristic χ_{KRH} is defined by

$$\chi_{KRH} \left(\bigoplus_{i,j \in \mathbb{Z}, k \in \mathbb{Z}_2} KRH^{i,j,k}(L) \right) := \sum_{i,j \in \mathbb{Z}, k \in \mathbb{Z}_2} (-1)^i \dim_{\mathbb{Q}}(KRH^{i,j,k}(L)) q^j.$$

This equals $\mathfrak{sl}_n$ quantum link invariant $\langle L \rangle_n$ for the link L .

$$\begin{aligned} \langle \text{crossing} \rangle_n &= q^{n-1} \langle \text{curved arrow} \rangle_n - q^n \langle \text{curved arrow} \rangle_n \\ \langle \text{crossing} \rangle_n &= q^{1-n} \langle \text{curved arrow} \rangle_n - q^{-n} \langle \text{curved arrow} \rangle_n \\ \langle D \sqcup D' \rangle_n &= \langle D \rangle_n \langle D' \rangle_n \quad \text{for any diagrams } D \text{ and } D' \\ \langle \text{circle} \rangle_n &= [n] \\ \langle \text{cup} \rangle_n &= [2] \langle \text{parallel lines} \rangle_n \\ \langle \text{cap} \rangle_n &= [n-1] \langle \text{curved arrow} \rangle_n \\ \langle \text{cup and cap} \rangle_n &= \langle \text{curved arrow} \rangle_n + [n-2] \langle \text{curved arrow} \rangle_n \\ \langle \text{crossing with cup and cap} \rangle_n + \langle \text{parallel lines} \rangle_n &= \langle \text{curved arrow} \rangle_n + \langle \text{parallel lines} \rangle_n \end{aligned}$$

where $[n]$ is a quantum integer $\frac{q^n - q^{-n}}{q - q^{-1}}$.

FIGURE 1. $\mathfrak{sl}_n$ quantum link invariant




FIGURE 2. Two type resolutions for crossing

In this paper, we define matrix factorizations for the more general local diagrams ,  and the double line . And we show that these matrix factorizations have some suitable properties; Proposition 3.5 and corollary 3.6 in Section 3.2.

Proposition 3.5 claims that the inner marking made by gluing these local diagrams can be removed in the homotopy category of a matrix factorization. For example, the matrix factorization for the diagram  is obtained as a tensor product of two matrix factorizations for  and .

M. Khovanov and L. Rozansky showed that the Euler characteristic of a matrix factorization for the single loop  equals to the quantum dimension of the fundamental representation V of $U_q(\mathfrak{sl}_n)$. Corollary 3.6 claims that the Euler characteristic of a matrix factorization for the double loop  also equals to the quantum dimension of $\wedge^2 V$ of $U_q(\mathfrak{sl}_n)$.

Acknowledgements: The author would like to thank Hiroaki Kanno, Hiroyuki Ochiai, Hidefumi Ohsugi and Akihiro Tsuchiya for many helpful discussion. And he would also like to thank Mikhail Khovanov and Lev Rozansky for supplying their paper [9] (preliminary version) to him.

2. CATEGORY OF MATRIX FACTORIZATION

2.1. Matrix factorization. We describe a category of a matrix factorization. M. Khovanov and L. Rozansky first imported this algebraic object into link theory [7][8].

Let R be a polynomial ring over $\mathbb{Q}$ and let M_0, M_1 be free R -modules permitted infinite rank. $\overline{M} = (M_0, M_1)$ is a *matrix factorization* with a potential $\omega \in R$ if $\overline{M}$ consists of a 2-cyclic complex of free R -modules M_0, M_1 ;

$$\left(M_0 \xrightarrow{d_{M_0}} M_1 \xrightarrow{d_{M_1}} M_0 \right)$$

such that $d_{M_1} d_{M_0} = \omega \text{Id}_{M_0}$ and $d_{M_0} d_{M_1} = \omega \text{Id}_{M_1}$.

For a polynomial ring R and a polynomial $\omega \in R$, let $\text{MF}_{R, \omega}$ be a category of a matrix factorization whose object is a matrix factorization (M_0, M_1) with the potential ω and whose morphism $\overline{f} = (f_0, f_1)$ between $\overline{M} = (M_0, M_1)$ and $\overline{N} = (N_0, N_1)$ consists of a pair of R -module morphisms $f_0 : M_0 \rightarrow N_0$ and $f_1 : M_1 \rightarrow N_1$ such that $d_{N_0} f_0 = f_1 d_{M_0}$ and $d_{N_1} f_1 = f_0 d_{M_1}$.

For $\overline{M} \in \text{Ob}(\text{MF}_{R, \omega})$ and $\overline{N} \in \text{Ob}(\text{MF}_{R, \omega})$, we define the tensor product $\overline{M} \boxtimes \overline{N} \in \text{Ob}(\text{MF}_{R \otimes R, \omega + \omega})$ by

$$\overline{M} \boxtimes \overline{N} := \left(\begin{pmatrix} M_0 \otimes N_0 \\ M_1 \otimes N_1 \end{pmatrix} \xrightarrow{\begin{pmatrix} d_{M_0} & -d_{N_1} \\ d_{N_0} & d_{M_1} \end{pmatrix}} \begin{pmatrix} M_1 \otimes N_0 \\ M_0 \otimes N_1 \end{pmatrix} \xrightarrow{\begin{pmatrix} d_{M_1} & d_{N_1} \\ -d_{N_0} & d_{M_0} \end{pmatrix}} \begin{pmatrix} M_0 \otimes N_0 \\ M_1 \otimes N_1 \end{pmatrix} \right),$$

where

$$\begin{pmatrix} d_{M_0} & -d_{N_1} \\ d_{N_0} & d_{M_1} \end{pmatrix} \text{ and } \begin{pmatrix} d_{M_1} & d_{N_1} \\ -d_{N_0} & d_{M_0} \end{pmatrix}$$

simply denote

$$\begin{pmatrix} d_{M_0} \otimes \text{Id}_{N_0} & -\text{Id}_{M_1} \otimes d_{N_1} \\ \text{Id}_{M_0} \otimes d_{N_0} & d_{M_1} \otimes \text{Id}_{N_1} \end{pmatrix} \text{ and } \begin{pmatrix} d_{M_1} \otimes \text{Id}_{N_0} & \text{Id}_{M_0} \otimes d_{N_1} \\ -\text{Id}_{M_1} \otimes d_{N_0} & d_{M_0} \otimes \text{Id}_{N_1} \end{pmatrix}.$$

Remark 2.1. We consider $R \otimes R'$ as a tensor product of polynomial rings R and R' over $R \cap R'$. We also consider $M \otimes M'$ as a tensor product of an R -module M and an R' -module M' over $R \cap R'$.

Lemma 2.2. (1) For $\overline{M} \in \text{Ob}(\text{MF}_{R,\omega})$ and $\overline{N} \in \text{Ob}(\text{MF}_{R,\omega'})$, there is an isomorphism in $\text{MF}_{R \otimes R', \omega + \omega'}$

$$\overline{M} \boxtimes \overline{N} \simeq \overline{N} \boxtimes \overline{M}.$$

(2) For $\overline{L} \in \text{Ob}(\text{MF}_{R,\omega})$, $\overline{M} \in \text{Ob}(\text{MF}_{R,\omega'})$ and $\overline{N} \in \text{Ob}(\text{MF}_{R',\omega''})$, there is an isomorphism in $\text{MF}_{R \otimes R \otimes R', \omega + \omega' + \omega''}$

$$(\overline{L} \boxtimes \overline{M}) \boxtimes \overline{N} \simeq \overline{L} \boxtimes (\overline{M} \boxtimes \overline{N}).$$

Proof. (1) For R -modules M and N , we define $T : M \otimes N \rightarrow N \otimes M$ by $T(m \otimes n) := n \otimes m$. $\overline{f} = \left(\begin{pmatrix} T & 0 \\ 0 & -T \end{pmatrix}, \begin{pmatrix} 0 & T \\ T & 0 \end{pmatrix} \right)$ is a morphism from $\overline{M} \boxtimes \overline{N}$ to $\overline{N} \boxtimes \overline{M}$ and also a morphism from $\overline{N} \boxtimes \overline{M}$ to $\overline{M} \boxtimes \overline{N}$. Since $\overline{f}^2 = \text{Id}$, $\overline{f}$ gives isomorphism between these matrix factorizations.

(2) By definition, we have

$$d_{((\overline{L} \boxtimes \overline{M}) \boxtimes \overline{N})_0} = \begin{pmatrix} d_{L_0} & -d_{M_1} & -d_{N_1} & 0 \\ d_{M_0} & d_{L_1} & 0 & -d_{N_1} \\ d_{N_0} & 0 & d_{L_1} & d_{M_1} \\ 0 & d_{N_0} & -d_{M_0} & d_{L_0} \end{pmatrix}, d_{((\overline{L} \boxtimes \overline{M}) \boxtimes \overline{N})_1} = \begin{pmatrix} d_{L_1} & d_{M_1} & d_{N_1} & 0 \\ -d_{M_0} & d_{L_0} & 0 & d_{N_1} \\ -d_{N_0} & 0 & d_{L_0} & -d_{M_1} \\ 0 & -d_{N_0} & d_{M_0} & d_{L_1} \end{pmatrix}$$

and

$$d_{(\overline{L} \boxtimes (\overline{M} \boxtimes \overline{N}))_0} = \begin{pmatrix} d_{L_0} & 0 & -d_{M_1} & -d_{N_1} \\ 0 & d_{L_0} & d_{N_0} & -d_{M_0} \\ d_{M_0} & -d_{N_1} & d_{L_1} & 0 \\ d_{N_0} & d_{M_1} & 0 & d_{L_1} \end{pmatrix}, d_{(\overline{L} \boxtimes (\overline{M} \boxtimes \overline{N}))_1} = \begin{pmatrix} d_{L_1} & 0 & d_{M_1} & d_{N_1} \\ 0 & d_{L_1} & -d_{N_0} & d_{M_0} \\ -d_{M_0} & d_{N_1} & d_{L_0} & 0 \\ -d_{N_0} & -d_{M_1} & 0 & d_{L_0} \end{pmatrix}.$$

Thus it is obvious that $\overline{f} = \left(\begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix}, \begin{pmatrix} 1 & 0 & 0 & 0 \\ 0 & 0 & 0 & 1 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{pmatrix} \right) : (\overline{L} \boxtimes \overline{M}) \boxtimes \overline{N} \rightarrow \overline{L} \boxtimes (\overline{M} \boxtimes \overline{N})$ is an isomorphism. $\square$

Lemma 2.3. The matrix factorization $(R \xrightarrow{0} 0 \xrightarrow{0} R)$ is the unit object in $\text{MF}_{R,\omega}$. That is, for any matrix factorization $\overline{M} \in \text{MF}_{R,\omega}$,

$$\overline{M} \boxtimes (R \xrightarrow{0} 0 \xrightarrow{0} R) \simeq \overline{M}.$$

Proof. By definition, we have

$$\begin{aligned} \overline{M} \boxtimes (R \xrightarrow{0} 0 \xrightarrow{0} R) &= \left(\begin{pmatrix} M_0 \otimes R \\ M_1 \otimes 0 \end{pmatrix} \xrightarrow{\begin{pmatrix} d_{M_0} & 0 \\ 0 & d_{M_1} \end{pmatrix}} \begin{pmatrix} M_1 \otimes R \\ M_0 \otimes 0 \end{pmatrix} \xrightarrow{\begin{pmatrix} d_{M_1} & 0 \\ 0 & d_{M_0} \end{pmatrix}} \begin{pmatrix} M_0 \otimes R \\ M_1 \otimes 0 \end{pmatrix} \right) \\ &\simeq \overline{M}. \end{aligned}$$

$\square$

The translation functor $\langle 1 \rangle$ changes the matrix factorization $\overline{M} = (M_0, M_1)$ into

$$\overline{M} \langle 1 \rangle = \left(M_1 \xrightarrow{-d_{M_1}} M_0 \xrightarrow{-d_{M_0}} M_1 \right).$$

The functor $\langle 2 \rangle (= \langle 1 \rangle^2)$ is the identity functor.

Lemma 2.4. *For $\overline{M} \in \text{Ob}(\text{MF}_{R,\omega})$ and $\overline{N} \in \text{Ob}(\text{MF}_{R,\omega'})$, there is an isomorphism in $\text{MF}_{R \otimes R, \omega + \omega'}$*

$$\begin{aligned} (\overline{M} \boxtimes \overline{N}) \langle 1 \rangle &= (\overline{M} \langle 1 \rangle) \boxtimes \overline{N} \\ &\simeq \overline{M} \boxtimes (\overline{N} \langle 1 \rangle). \end{aligned}$$

Proof. We directly find that $(\overline{M} \boxtimes \overline{N}) \langle 1 \rangle$ equals $(\overline{M} \langle 1 \rangle) \boxtimes \overline{N}$ by definition. The second equivalence is correct by Lemma 2.2 (1) and the first equality. $\square$

The morphism $\overline{f} : \overline{M} \rightarrow \overline{N} \in \text{Mor}(\text{MF}_{R,\omega})$ is *null-homotopic* if morphisms $h_0 : M_0 \rightarrow N_0$ and $h_1 : M_1 \rightarrow N_1$ exist such that $f_0 = h_1 d_{M_0} + d_{N_1} h_0$ and $f_1 = h_0 d_{M_1} + d_{N_0} h_1$. And $\overline{f}, \overline{g} : \overline{M} \rightarrow \overline{N} \in \text{Mor}(\text{MF}_{R,\omega})$ are *homotopic* if $\overline{f} - \overline{g}$ is null-homotopic.

Let $\text{HMF}_{R,\omega}$ be the quotient category of $\text{MF}_{R,\omega}$ which has the same objects to $\text{Ob}(\text{MF}_{R,\omega})$ and has morphisms of $\text{Mor}(\text{MF}_{R,\omega})$ modulo null-homotopic. A matrix factorization in $\text{MF}_{R,\omega}$ is called *contractible* if it is isomorphic to the zero matrix factorization

$$\left(0 \xrightarrow{0} 0 \xrightarrow{0} 0 \right)$$

in $\text{HMF}_{R,\omega}$.

Example 2.5. *Let R_0, R_1 be a ring R and $\omega \in R$.*

$$\left(R_1 \xrightarrow{\omega} R_0 \xrightarrow{1} R_1 \right)$$

and

$$\left(R_1 \xrightarrow{1} R_0 \xrightarrow{\omega} R_1 \right)$$

are contractible.

2.2. $\mathbb{Z}$ -graded matrix factorization. Let R be replaced with a $\mathbb{Z}$ -graded polynomial ring over $\mathbb{Q}$ whose each parameter has a $\mathbb{Z}$ -grading and let M_0, M_1 be also replaced with free $\mathbb{Z}$ -graded R -modules. The category $\text{MF}_{R,\omega}^{gr}$ is the category of a $\mathbb{Z}$ -graded matrix factorization whose object is the same object to $\text{Ob}(\text{MF}_{R,\omega})$ except having a $\mathbb{Z}$ -grading and whose morphism consists of a morphism with preserving a $\mathbb{Z}$ -grading in $\text{Mor}(\text{MF}_{R,\omega})$. The quotient category $\text{HMF}_{R,\omega}^{gr}$ is also defined as $\text{Ob}(\text{HMF}_{R,\omega}^{gr}) = \text{Ob}(\text{MF}_{R,\omega}^{gr})$ and $\text{Mor}(\text{HMF}_{R,\omega}^{gr}) = \text{Mor}(\text{MF}_{R,\omega}^{gr}) / \{ \text{null-homotopic} \}$.

The $\mathbb{Z}$ -grading shift $\{n\}$ ($n \in \mathbb{Z}$) turns the matrix factorization $\overline{M}$ into

$$\left(M_0 \{n\} \xrightarrow{d_{M_0}} M_1 \{n\} \xrightarrow{d_{M_1}} M_0 \{n\} \right).$$

Lemma 2.6. *For $\overline{M} \in \text{Ob}(\text{MF}_{R,\omega}^{gr})$ and $\overline{N} \in \text{Ob}(\text{MF}_{R,\omega'}^{gr})$, there is an equality in $\text{MF}_{R \otimes R, \omega + \omega'}^{gr}$*

$$\begin{aligned} (\overline{M} \boxtimes \overline{N}) \{n\} &= (\overline{M} \{n\}) \boxtimes \overline{N} \\ &= \overline{M} \boxtimes (\overline{N} \{n\}) \end{aligned}$$

Proof. We find that these objects are really identical by definition. $\square$

For the $\mathbb{Z}$ -graded matrix factorization $\overline{M}$ in $\text{MF}_{R,0}^{gr}$, we can consider $\mathbb{Z} \oplus \mathbb{Z}_2$ -graded homology group as follows;

$$H(\overline{M}) = \bigoplus_{j \in \mathbb{Z}, k \in \mathbb{Z}_2} H^{j,k}(\overline{M})$$

whose k is a complex grading of the matrix factorization $\overline{M}$, i.e. $i = 0$ or 1 , and j is a $\mathbb{Z}$ -grading induced by the $\mathbb{Z}$ -graded modules of the matrix factorization $\overline{M}$. The *Euler characteristic* $\overline{\chi}$ is defined by

$$\overline{\chi}(\mathrm{H}(\overline{M})) = \sum_{j \in \mathbb{Z}, k \in \mathbb{Z}_2} \dim_{\mathbb{Q}} \mathrm{H}^{j,k}(\overline{M}) q^j.$$

2.3. Koszul matrix factorization. Let R be a $\mathbb{Z}$ -graded polynomial ring over $\mathbb{Q}$ and let $\deg_{\mathbb{Z}}(a)$ denote a $\mathbb{Z}$ -grading of the polynomial $a \in R$. For polynomials $a, b \in R$ and a $\mathbb{Z}$ -graded R -module M , we define the matrix factorization $K(a; b)_M$ with the potential ab by

$$K(a; b)_M := \left(M \xrightarrow{a} M\{\tfrac{1}{2}(\deg_{\mathbb{Z}}(b) - \deg_{\mathbb{Z}}(a))\} \xrightarrow{b} M \right).$$

Remark 2.7. Let R be a polynomial ring over $\mathbb{Q}$ and let R_y be the polynomial ring $R[y]$. For polynomials a and b in R , $K(a; b)_{R_y}$ is a matrix factorization of R_y -modules with rank 1 as an object in $\mathrm{MF}_{R_y, ab}^{gr}$. And we can consider that $K(a; b)_{R_y}$ is a matrix factorization of R -modules with infinite rank as an object in $\mathrm{MF}_{R, ab}^{gr}$.

Lemma 2.8.

$$K(a; b)_M \langle 1 \rangle = K(-b; -a)_M \left\{ \tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a) \right\}.$$

Proof. By definition, we have

$$\begin{aligned} K(a; b)_M \langle 1 \rangle &= \left(M \xrightarrow{a} M\{\tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a)\} \xrightarrow{b} M \right) \langle 1 \rangle \\ &= \left(M\{\tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a)\} \xrightarrow{-b} M \xrightarrow{-a} M\{\tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a)\} \right) \\ &= \left(M \xrightarrow{-b} M\{\tfrac{1}{2}(\deg_{\mathbb{Z}} a - \deg_{\mathbb{Z}} b)\} \xrightarrow{-a} M \right) \left\{ \tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a) \right\} \\ &= K(-b; -a)_M \left\{ \tfrac{1}{2}(\deg_{\mathbb{Z}} b - \deg_{\mathbb{Z}} a) \right\}. \end{aligned}$$

□

In general, for the sequences $\mathbf{a} = {}^t(a_1, a_2, \dots, a_k)$ and $\mathbf{b} = {}^t(b_1, b_2, \dots, b_k)$ of polynomials in R , we define the matrix factorization $K(\mathbf{a}; \mathbf{b})_M$ with the potential $\sum_{i=1}^k a_i b_i$ by

$$K(\mathbf{a}; \mathbf{b})_M := \bigotimes_{i=1}^k K(a_i; b_i)_M.$$

This matrix factorization is called a *Koszul matrix factorization*.

Lemma 2.9. Let c be a non-zero element in $\mathbb{Q}$.

There is an isomorphism in $\mathrm{MF}_{R, ab}^{gr}$

$$K(a; b)_M \simeq K(ca; c^{-1}b)_M.$$

Proof. $\overline{f} = (1, c) : K(a; b)_M \rightarrow K(ca; c^{-1}b)_M$ and $\overline{g} = (1, c^{-1}) : K(ca; c^{-1}b)_M \rightarrow K(a; b)_M$ satisfy that $\overline{g}\overline{f} = \overline{\mathrm{Id}}$ and $\overline{f}\overline{g} = \overline{\mathrm{Id}}$. Thus, these matrix factorizations are isomorphic.

□

Theorem 2.10. [Khovanov-Rozansky, Theorem 2.2.[9]] We put $R = \mathbb{Q}[x_1, x_2, \dots, x_n]$ and $R_y = R[y]$. Let ${}^t(a_1, a_2, \dots, a_k)$ and ${}^t(b_1, b_2, \dots, b_k)$ be sequences of polynomials $a_j, b_j \in R_y$.

We assume that $K(\mathbf{a}; \mathbf{b})_{R_y}$ is a Koszul matrix factorization with the potential $\omega \in R$. That is to say, we can see that this matrix factorization is an object in $\mathrm{HMF}_{R, \omega}^{gr}$, which consists of infinite rank R -modules R_y .

Furthermore, we assume that $b_i = cy^n + p$ for some i , where c is a non-zero element in $\mathbb{Q}$ and p is the polynomial in R_y whose degree for y is less than n . Then, there is an isomorphism in $\text{HMF}_{R,\omega}^{gr}$

$$K(\mathbf{a}; \mathbf{b})_{R_y} \simeq K(\overset{i}{\mathbf{a}}; \overset{i}{\mathbf{b}})_{R_y / \langle b_i \rangle},$$

where $\overset{i}{\mathbf{a}}$ and $\overset{i}{\mathbf{b}}$ are associated with $\mathbf{a}$ and $\mathbf{b}$ removing the i -th polynomial.

Proof. We can replace b_i with $y^n + p$ using Lemma 2.9. Then, we repeat proof by M.Khovanov and L.Rozansky in [9]. By definition and the above assumption, we have

$$K(\mathbf{a}; \mathbf{b})_{R_y} = K(\overset{i}{\mathbf{a}}; \overset{i}{\mathbf{b}})_{R_y} \boxtimes K(a_i; y^n + p)_{R_y}.$$

The Koszul matrix factorization $K(\overset{i}{\mathbf{a}}; \overset{i}{\mathbf{b}})_{R_y}$ is described as

$$\left(R_y^r \xrightarrow{D_0} R_y^r \xrightarrow{D_1} R_y^r \right),$$

where $D_0, D_1 \in M_r(R_y)$ and $r = 2^{k-2}$. Thus, we obtain

$$K(\mathbf{a}; \mathbf{b})_{R_y} = \left(\begin{array}{c} R_y^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{D_0'} \begin{array}{c} R_y^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{D_1'} \begin{array}{c} R_y^r \\ \oplus \\ R_y^r \end{array} \right),$$

$$\text{where } D_0' = \begin{pmatrix} D_0 & (-y^n - p)\text{Id}_{R_y^r} \\ a_i \text{Id}_{R_y^r} & D_1 \end{pmatrix}, \quad D_1' = \begin{pmatrix} D_1 & (y^n + p)\text{Id}_{R_y^r} \\ -a_i \text{Id}_{R_y^r} & D_0 \end{pmatrix}.$$

The ring R_y is split into the direct sum as an R -module as follows;

$$R_y \simeq R_{<n} \oplus R_{\geq n}, \text{ where } R_{<n} = \bigoplus_{i=0}^{n-1} R \cdot y^i \text{ and } R_{\geq n} = \bigoplus_{i=0}^{\infty} R \cdot y^i (y^n + p).$$

The R -module morphism $R_y \xrightarrow{y^n+p} R_y$ induces the R -module isomorphism

$$f_{iso} : R_y \xrightarrow{y^n+p} R_{\geq n}.$$

Moreover, there are the natural R -module injection

$$f_{inj} : R_{<n} \longrightarrow R_y,$$

and the natural R -module projections

$$f_{proj_{<n}} : R_y \longrightarrow R_{<n}$$

and

$$f_{proj_{\geq n}} : R_y \longrightarrow R_{\geq n}.$$

The R -module R_y^r is also split into the direct sum

$$R_y^r \simeq R_{<n}^r \oplus R_{\geq n}^r.$$

Then, there are the R -module isomorphism

$$F_{iso} : R_y^r \xrightarrow{(y^n+p)\text{Id}_{R_y^r}} R_{\geq n}^r,$$

the R -module injection

$$F_{inj} : R_{<n}^r \longrightarrow R_y^r$$

and the R -module projections

$$F_{proj<_n} : R_y^r \longrightarrow R_{<_n}^r$$

and

$$F_{proj\geq_n} : R_y^r \longrightarrow R_{\geq_n}^r.$$

It is easy to find that the R -module morphisms

$$\phi_0 = \begin{array}{c} R_{<_n}^r \\ \oplus \\ R_{\geq_n}^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{\begin{pmatrix} F_{inj} & (y^n + p)F_{iso}^{-1} & 0 \\ 0 & D_0 F_{iso}^{-1} & \text{Id}_{R_y^r} \end{pmatrix}} \begin{array}{c} R_y^r \\ \oplus \\ R_y^r \end{array}$$

and

$$\phi_1 = \begin{array}{c} R_{<_n}^r \\ \oplus \\ R_{\geq_n}^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{\begin{pmatrix} F_{inj} & (-y^n - p)F_{iso}^{-1} & 0 \\ 0 & D_1 F_{iso}^{-1} & \text{Id}_{R_y^r} \end{pmatrix}} \begin{array}{c} R_y^r \\ \oplus \\ R_y^r \end{array}$$

are R -module isomorphisms. Let $\overline{M}$ be the matrix factorization

$$\overline{M} = \left(\begin{array}{c} R_{<_n}^r \\ \oplus \\ R_{\geq_n}^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{\phi_1^{-1} D_0' \phi_0} \begin{array}{c} R_{<_n}^r \\ \oplus \\ R_{\geq_n}^r \\ \oplus \\ R_y^r \end{array} \xrightarrow{\phi_0^{-1} D_1' \phi_1} \begin{array}{c} R_{<_n}^r \\ \oplus \\ R_{\geq_n}^r \\ \oplus \\ R_y^r \end{array} \right).$$

Then, $\overline{\phi} = (\phi_0, \phi_1)$ is an isomorphism from $\overline{M}$ to $K(\mathbf{a}; \mathbf{b})_{R_y}$ in $\text{MF}_{R, \omega}^{gr}$.

Since $\overline{\phi}^{-1}$ consists of

$$\phi_0^{-1} = \begin{pmatrix} F_{proj<_n} & 0 \\ (y^n + p)F_{iso}^{-1} F_{proj\geq_n} & 0 \\ -D_0 F_{iso}^{-1} F_{proj\geq_n} & \text{Id}_{R_y^r} \end{pmatrix} \text{ and } \phi_1^{-1} = \begin{pmatrix} F_{proj<_n} & 0 \\ (-y^n - p)F_{iso}^{-1} F_{proj\geq_n} & 0 \\ D_1 F_{iso}^{-1} F_{proj\geq_n} & \text{Id}_{R_y^r} \end{pmatrix},$$

the morphisms $\phi_1^{-1} D_0' \phi_0$ and $\phi_0^{-1} D_1' \phi_1$ are described by

$$\phi_1^{-1} D_0' \phi_0 = \begin{pmatrix} F_{proj<_n} D_0 F_{inj} & 0 & 0 \\ * & 0 & F_{iso} \\ * & \omega F_{iso}^{-1} & 0 \end{pmatrix} \text{ and } \phi_0^{-1} D_1' \phi_1 = \begin{pmatrix} F_{proj<_n} D_1 F_{inj} & 0 & 0 \\ * & 0 & F_{iso} \\ * & \omega F_{iso}^{-1} & 0 \end{pmatrix}.$$

The matrix factorization obtained by restricting $R_{<_n}^r \oplus R_{\geq_n}^r \oplus R_y^r$ of $\overline{M}$ to $R_{\geq_n}^r \oplus R_y^r$

$$\left(\begin{pmatrix} R_{\geq_n}^r \\ \oplus \\ R_y^r \end{pmatrix} \xrightarrow{\begin{pmatrix} 0 & F_{iso} \\ \omega F_{iso}^{-1} & 0 \end{pmatrix}} \begin{pmatrix} R_{\geq_n}^r \\ \oplus \\ R_y^r \end{pmatrix} \xrightarrow{\begin{pmatrix} 0 & F_{iso} \\ \omega F_{iso}^{-1} & 0 \end{pmatrix}} \begin{pmatrix} R_{\geq_n}^r \\ \oplus \\ R_y^r \end{pmatrix} \right)$$

is contractible in $\text{HMF}_{R, \omega}^{gr}$. Thus, $\overline{M}$ is isomorphic to the quotient matrix factorization

$$\left(R_{<_n}^r \xrightarrow{F_{proj<_n} D_0 F_{inj}} R_{<_n}^r \xrightarrow{F_{proj<_n} D_1 F_{inj}} R_{<_n}^r \right)$$

in $\text{HMF}_{R,\omega}^{gr}$. By the choice of a basis of R_y as an R -module, it is easy to find that this quotient matrix factorization equals $K(\overset{i}{\mathbf{a}}; \overset{i}{\mathbf{b}})_{R_y/\langle b_i \rangle}$. $\square$

3. MATRIX FACTORIZATION FOR TRIVALENT DIAGRAMS AND DOUBLE LINE

3.1. Khovanov Rozansky homology. We briefly recall Khovanov-Rozansky link homology theory, only the definition of matrix factorizations for planar diagrams and some properties of the matrix factorizations. See Section 6 in [7] for further details.

In [7], M. Khovanov and L. Rozansky defined a categorification of planar graphs $\begin{array}{c} \leftarrow \\ \text{---} \\ \rightarrow \end{array}$ and $\curvearrowright$ as a Koszul matrix factorization.

We assign a parameter x_i on the end point of a single line. And each $\mathbb{Z}$ -grading of parameters x_i is 2. This $\mathbb{Z}$ -grading induces a $\mathbb{Z}$ -grading of a matrix factorization in $\text{MF}_{R,\omega}^{gr}$ and $\text{HMF}_{R,\omega}^{gr}$. The function $f(s_1, s_2)$ is obtained by expanding the power sum $x^{n+1} + y^{n+1}$ with the elementary symmetric polynomials $x + y$ and xy , i.e. the function $f(s_1, s_2)$ satisfies that

$$f(x + y, xy) = x^{n+1} + y^{n+1}.$$

We define matrix factorizations for the planar diagrams $\begin{array}{c} x_1 \\ \curvearrowright \\ x_2 \end{array}$ and $\begin{array}{c} x_1 \leftarrow x_2 \\ \text{---} \\ x_3 \rightarrow x_4 \end{array}$ by

$$\begin{aligned} \mathcal{C}\left(\begin{array}{c} x_1 \\ \curvearrowright \\ x_2 \end{array}\right)_n &= K(\pi_{12}; x_1 - x_2)_{\mathbb{Q}[x_1, x_2]}, \\ \mathcal{C}\left(\begin{array}{c} x_1 \leftarrow x_2 \\ \text{---} \\ x_3 \rightarrow x_4 \end{array}\right)_n &= K\left(\left(\begin{array}{c} u_{3,4}^{1,2} \\ v_{3,4}^{1,2} \end{array}\right); \left(\begin{array}{c} x_1 + x_2 - x_3 - x_4 \\ x_1 x_2 - x_3 x_4 \end{array}\right)\right)_{\mathbb{Q}[x_1, x_2, x_3, x_4]} \{-1\}, \end{aligned}$$

where $\pi_{12}(x_1 - x_2) = x_1^{n+1} - x_2^{n+1}$, $u_{3,4}^{1,2} = \frac{f(x_1+x_2, x_1 x_2) - f(x_3+x_4, x_1 x_2)}{x_1+x_2-x_3-x_4}$ and $v_{3,4}^{1,2} = \frac{f(x_3+x_4, x_1 x_2) - f(x_3+x_4, x_3 x_4)}{x_1 x_2 - x_3 x_4}$.

Furthermore, we construct a matrix factorization for the more complex planar diagrams produced by combinatorially joining the diagrams $\begin{array}{c} \leftarrow \\ \text{---} \\ \rightarrow \end{array}$ and $\curvearrowright$. Now, suppose the planar diagrams $\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}$ and $\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}$, which can match at x and x' with keeping the orientation. For the matrix factorizations

$$\mathcal{C}\left(\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}\right)_n \in \text{Ob}(\text{MF}_{R,\omega+x^{n+1}}^{gr}) \quad \text{and} \quad \mathcal{C}\left(\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}\right)_n \in \text{Ob}(\text{MF}_{R,\omega'-x^{n+1}}^{gr}),$$

we define the matrix factorization

$$\mathcal{C}\left(\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}\right)_n \in \text{Ob}(\text{MF}_{R \otimes R, \omega+\omega'}^{gr}) \quad \text{by} \quad \mathcal{C}\left(\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}\right)_n \boxtimes \mathcal{C}\left(\begin{array}{c} \curvearrowright \\ \text{---} \\ \curvearrowright \end{array}\right)_n \Big|_{x=x'}.$$

This means that we identify the top parameter x and the tail parameter x' after taking the tensor product of these matrix factorizations. And if a planar diagram consists of two disjoint planar diagrams D_1 and D_2 , then this matrix factorization is defined by the tensor product of matrix factorizations for D_1 and D_2 ;

$$\mathcal{C}(D_1 \sqcup D_2)_n = \mathcal{C}(D_1)_n \boxtimes \mathcal{C}(D_2)_n.$$

Proposition 3.1. *The map $\mathcal{C}$ from a planar diagram to a matrix factorization is well-defined.*

Proof. This is obvious by Lemma 2.2. $\square$

The next Proposition from (1) to (7) claims that the inner marking built by gluing diagrams can be removed.

Proposition 3.2. (1) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x] \otimes R, x^{n+1}-\omega}^{gr}$, where R is a polynomial ring generated by boundary parameters except x ,*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with an incoming arrow from the left labeled } x, \text{ and an outgoing arrow to the right labeled } x. \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with an incoming arrow from the left labeled } x, \text{ and an outgoing arrow to the right labeled } x. \end{array} \right)_n.$$

(2) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x] \otimes R, \omega-x^{n+1}}^{gr}$, where R is a polynomial ring generated by boundary parameters except x ,*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with an incoming arrow from the left labeled } x, \text{ and an outgoing arrow to the right labeled } x'. \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with an incoming arrow from the left labeled } x, \text{ and an outgoing arrow to the right labeled } x. \end{array} \right)_n.$$

(3) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}, 0}^{gr}$*

$$\begin{aligned} \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with an incoming arrow from the left labeled } x_1. \end{array} \right)_n &= K(\pi_{12}; x_1 - x_2)_{\mathbb{Q}[x_1, x_2]} \Big|_{x_1=x_2} \\ &\simeq (0 \rightarrow \mathbb{Q}[x] / \langle x^n \rangle \{1 - n\} \rightarrow 0). \end{aligned}$$

(4) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x_1, x_2, x_5, x_6], x_1^{n+1}+x_2^{n+1}-x_5^{n+1}-x_6^{n+1}}^{gr}$*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with four incoming arrows from the left labeled } x_1, x_2, x_3, x_4 \text{ and two outgoing arrows to the right labeled } x_5, x_6. \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and two outgoing arrows to the right labeled } x_5, x_6. \end{array} \right)_n \{-1\} \oplus \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and two outgoing arrows to the right labeled } x_5, x_6. \end{array} \right)_n \{1\}.$$

(5) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x_1, x_2], x_1^{n+1}-x_2^{n+1}}^{gr}$*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and one outgoing arrow to the right labeled } x_3. \end{array} \right)_n \simeq \bigoplus_{i=0}^{n-2} \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with one incoming arrow from the left labeled } x_1 \text{ and one outgoing arrow to the right labeled } x_2. \end{array} \right)_n \{2 - n + 2i\} \langle 1 \rangle.$$

(6) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x_1, x_2, x_3, x_4], x_1^{n+1}-x_2^{n+1}+x_3^{n+1}-x_4^{n+1}}^{gr}$*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with four incoming arrows from the left labeled } x_1, x_2, x_3, x_4 \text{ and two outgoing arrows to the right labeled } x_5, x_6. \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and two outgoing arrows to the right labeled } x_3, x_4. \end{array} \right)_n \oplus \bigoplus_{i=0}^{n-3} \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with one incoming arrow from the left labeled } x_1 \text{ and one outgoing arrow to the right labeled } x_2. \end{array} \right)_n \{3 - n + 2i\}.$$

(7) *There is an isomorphism in $\mathrm{HMF}_{\mathbb{Q}[x_1, x_2, x_3, x_4, x_5, x_6], x_1^{n+1}+x_2^{n+1}+x_3^{n+1}-x_4^{n+1}-x_5^{n+1}-x_6^{n+1}}^{gr}$*

$$\mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with six incoming arrows from the left labeled } x_1, x_2, x_3, x_4, x_5, x_6. \end{array} \right)_n \oplus \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and one outgoing arrow to the right labeled } x_3. \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with six incoming arrows from the left labeled } x_1, x_2, x_3, x_4, x_5, x_6. \end{array} \right)_n \oplus \mathcal{C}' \left(\begin{array}{c} \text{Diagram: a circle with two incoming arrows from the left labeled } x_1, x_2 \text{ and one outgoing arrow to the right labeled } x_3. \end{array} \right)_n.$$

Proof. See [7]. □

The matrix factorization for the single loop $\bigcirc$ is defined as the following matrix factorization;

$$\mathcal{C}(\bigcirc) = \left(0 \rightarrow \mathbb{Q}[x]/\langle x^n \rangle \{1-n\} \rightarrow 0 \right).$$

Since the potential of the matrix factorization $\mathcal{C}(\bigcirc)_n$ is 0, we can take the homology group H of this matrix factorization.

Corollary 3.3. *The Euler characteristic of the homology $H(\mathcal{C}(\bigcirc)_n)$ equals the value of $\mathfrak{sl}_n$ quantum link invariant for the single loop;*

$$\overline{\chi}(H(\mathcal{C}(\bigcirc)_n)) = [n].$$

Proof. See [7]. □

The map $\mathcal{C}$ from a planar diagram to a matrix factorization extends the map $\mathcal{C}$ from the projection diagram associated with a link to a complex of matrix factorizations.

$$\begin{aligned} \mathcal{C}(\nearrow \searrow) &:= \left(\longrightarrow \mathcal{C}(\begin{array}{c} \xrightarrow{-1} \\ \text{---} \\ \xrightarrow{-1} \end{array}) \{n\} \langle 1 \rangle \longrightarrow \mathcal{C}(\begin{array}{c} \xrightarrow{0} \\ \text{---} \\ \xrightarrow{0} \end{array}) \{n-1\} \langle 1 \rangle \longrightarrow 0 \longrightarrow \right) \\ \mathcal{C}(\nwarrow \nearrow) &:= \left(\longrightarrow 0 \longrightarrow \mathcal{C}(\begin{array}{c} \xrightarrow{0} \\ \text{---} \\ \xrightarrow{0} \end{array}) \{-n+1\} \langle 1 \rangle \longrightarrow \mathcal{C}(\begin{array}{c} \xrightarrow{1} \\ \text{---} \\ \xrightarrow{1} \end{array}) \{-n\} \langle 1 \rangle \longrightarrow \right) \end{aligned}$$

Theorem 3.4. [Khovanov-Rozansky,[7]] *The map $\mathcal{C}$ is an oriented link invariant in the homotopy category of a complex of matrix factorizations.*

Proof. See [7]. □

3.2. Definition of matrix factorization for trivalent diagrams and double line. We extend the map $\mathcal{C}$ for a planar diagram with two kinds of the boundaries, which consist of end points of a single line and a double line. We define matrix factorizations for the essential planar diagrams $\begin{array}{c} \nearrow \\ \nwarrow \end{array}$, $\begin{array}{c} \uparrow \\ \downarrow \end{array}$ and $\begin{array}{c} \nearrow \\ \nwarrow \end{array}$ as follows.

After this, we assume that $n \geq 3$. We assign the parameter x_i on the end point of a single line and the pair of parameters (y_j, z_j) on the end point of a double line. And each $\mathbb{Z}$ -grading of x_i and y_j also equals 2, but the $\mathbb{Z}$ -grading of z_i equals 4;

$$\deg_{\mathbb{Z}}(x_i) = 2, \deg_{\mathbb{Z}}(y_i) = 2 \text{ and } \deg_{\mathbb{Z}}(z_i) = 4.$$

Define a matrix factorization for the first diagram, double line, by

$$\mathcal{C}\left(\begin{array}{c} (y_1, z_1) \\ \nearrow \searrow \\ (y_2, z_2) \end{array}\right)_n = K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_2, z_1)}{y_1 - y_2} \\ \frac{f(y_2, z_1) - f(y_2, z_2)}{z_1 - z_2} \end{array} \right); \left(\begin{array}{c} y_1 - y_2 \\ z_1 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, y_2, z_1, z_2]}$$

in $\text{MF}^{gr}_{\mathbb{Q}[y_1, y_2, z_1, z_2], f(y_1, z_1) - f(y_2, z_2)}$.

Define a matrix factorization for the second trivalent diagram by

$$\mathcal{C}\left(\begin{array}{c} (y_3, z_3) \\ \uparrow \\ x_1 \nwarrow x_2 \end{array}\right)_n = K \left(\left(\begin{array}{c} \frac{f(y_3, z_3) - f(x_1 + x_2, z_3)}{y_3 - x_1 - x_2} \\ \frac{f(x_1 + x_2, z_3) - f(x_1 + x_2, x_1 x_2)}{z_3 - x_1 x_2} \end{array} \right); \left(\begin{array}{c} y_3 - x_1 - x_2 \\ z_3 - x_1 x_2 \end{array} \right) \right)_{\mathbb{Q}[x_1, x_2, y_3, z_3]}$$

in $\text{MF}_{\mathbb{Q}[x_1, x_2, y_3, z_3], f(y_3, z_3) - x_1^{n+1} - x_2^{n+1}}^{gr}$.

Define a matrix factorization for the third trivalent diagram by

$$\mathcal{C}' \left(\begin{array}{c} x_1 \swarrow \quad \searrow x_2 \\ \uparrow \\ (y_3, z_3) \end{array} \right)_n = K \left(\left(\begin{array}{c} \frac{f(x_1+x_2, x_1x_2) - f(y_3, x_1x_2)}{x_1+x_2-y_3} \\ \frac{f(y_3, x_1x_2) - f(y_3, z_3)}{x_1x_2-z_3} \end{array} \right); \begin{pmatrix} x_1+x_2-y_3 \\ x_1x_2-z_3 \end{pmatrix} \right)_{\mathbb{Q}[x_1, x_2, y_3, z_3]} \{-1\}$$

in $\text{MF}_{\mathbb{Q}[x_1, x_2, y_3, z_3], x_1^{n+1} + x_2^{n+1} - f(y_3, z_3)}^{gr}$.

The basic potential of a double line is the polynomial $f(y, z)$ obtained by expanding the power sum with the elementary symmetric polynomials. This polynomial $f(y, z)$ is a non-homogeneous polynomial, but it has the homogeneous $\mathbb{Z}$ -grading $2n+2$.

Now, we consider two planar diagrams, which can match at end points of oriented double lines with keeping the orientation. For the matrix factorizations $\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \in \text{Ob}(\text{MF}_{R, \omega + f(y, z)}^{gr})$ and $\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \in \text{Ob}(\text{MF}_{R, \omega - f(y, z)}^{gr})$, the matrix factorization $\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n$ is defined by the tensor product

$$\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \boxtimes \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \Big|_{(y, z) = (y, z)} \in \text{Ob}(\text{MF}_{R \otimes R, \omega + \omega}^{gr}).$$

For a polynomial $f \in \mathbb{Q}[x_1, x_2, \dots, x_k]$, the Jacobi algebra J_f is defined as the quotient ring

$$\mathbb{Q}[x_1, x_2, \dots, x_k] / \left\langle \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_k} \right\rangle,$$

where $\left\langle \frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_k} \right\rangle$ is the ideal of $\mathbb{Q}[x_1, x_2, \dots, x_k]$ generated by $\frac{\partial f}{\partial x_1}, \frac{\partial f}{\partial x_2}, \dots, \frac{\partial f}{\partial x_k}$.

Proposition 3.5. (1) *There is an isomorphism in $\text{HMF}_{\mathbb{Q}[y, z] \otimes R, f(y, z) - \omega}^{gr}$, where R is a polynomial ring generated by boundary parameters except y and z ,*

$$\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n.$$

(2) *There is an isomorphism in $\text{HMF}_{\mathbb{Q}[y, z] \otimes R, \omega - f(y, z)}^{gr}$, where R is a polynomial ring generated by boundary parameters except y and z ,*

$$\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y, z) \end{array} \right)_n.$$

(3) *There is an isomorphism in $\text{HMF}_{\mathbb{Q}, 0}^{gr}$*

$$\begin{aligned} \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y_1, z_1) \end{array} \right)_n &= K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_2, z_1)}{y_1 - y_2} \\ \frac{f(y_2, z_1) - f(y_2, z_2)}{z_1 - z_2} \end{array} \right); \begin{pmatrix} y_1 - y_2 \\ z_1 - z_2 \end{pmatrix} \right)_{\mathbb{Q}[y_1, y_2, z_1, z_2]} \Big|_{(y_1, z_1) = (y_2, z_2)} \\ &\simeq \left(J_{f(y, z)} \{4 - 2n\} \longrightarrow 0 \longrightarrow J_{f(y, z)} \{4 - 2n\} \right). \end{aligned}$$

(4) *There is an isomorphism in $\text{HMF}_{\mathbb{Q}[y_1, y_2, z_1, z_2], f(y_1, z_1) - f(y_2, z_2)}^{gr}$*

$$\mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y_1, z_1), (y_2, z_2) \end{array} \right)_n \simeq \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y_2, z_2) \end{array} \right)_n \{-1\} \oplus \mathcal{C}' \left(\begin{array}{c} \text{diagram} \\ (y_2, z_2) \end{array} \right)_n \{1\}.$$

(5) There is an isomorphism in $\text{HMF}^{gr}_{\mathbb{Q}[x_1, x_2, x_3, x_4], x_1^{n+1} + x_2^{n+1} - x_3^{n+1} - x_4^{n+1}}$

$$\mathcal{C} \left(\begin{array}{c} x_1 \leftarrow x_2 \\ \text{---} (y_5, z_5) \text{---} \\ x_3 \rightarrow x_4 \end{array} \right)_n \simeq \mathcal{C} \left(\begin{array}{c} x_1 \leftarrow x_2 \\ \text{---} \\ x_3 \rightarrow x_4 \end{array} \right)_n.$$

Proof. (1) The matrix factorization $\mathcal{C} \left(\begin{array}{c} (y', z') \\ \text{---} \\ (y, z) \end{array} \right)_n$ is described by

$$(M_0 \longrightarrow M_1 \longrightarrow M_0),$$

where M_i is an $R \otimes \mathbb{Q}[x', y']$ -module. Then, we have

$$\mathcal{C} \left(\begin{array}{c} (y', z') \\ \text{---} \\ (y, z) \end{array} \right)_n = (M_0 \longrightarrow M_1 \longrightarrow M_0) \boxtimes K \left(\left(\begin{array}{c} \frac{f(y, z) - f(y', z)}{y - y'} \\ \frac{f(y, z) - f(y', z)}{z - z'} \end{array} \right); \left(\begin{array}{c} y - y' \\ z - z' \end{array} \right) \right)_{\mathbb{Q}[y, y', z, z']}$$

Since the potential of this matrix factorization does not include the parameters y' and z' , we can apply Theorem 2.10 to $y - y'$ and $z - z'$. Hence, we obtain the following isomorphism

$$\begin{aligned} & (M_0 \longrightarrow M_1 \longrightarrow M_0) \boxtimes K \left(\left(\begin{array}{c} \frac{f(y, z) - f(y', z)}{y - y'} \\ \frac{f(y, z) - f(y', z)}{z - z'} \end{array} \right); \left(\begin{array}{c} y - y' \\ z - z' \end{array} \right) \right)_{\mathbb{Q}[y, y', z, z]} \\ & \simeq (M_0 / \langle y - y', z - z' \rangle M_0 \longrightarrow M_1 / \langle y - y', z - z' \rangle M_1 \longrightarrow M_0 / \langle y - y', z - z' \rangle M_0) \\ & \simeq \mathcal{C} \left(\begin{array}{c} \text{---} \\ (y, z) \end{array} \right)_n. \end{aligned}$$

(2) This proof is similar to (1)

(3) The polynomial $f(y_1, z_1)$ is explicitly described as

$$f(y_1, z_1) = y_1^{n+1} + (n+1) \sum_{1 \leq 2i \leq n+1} \frac{(-1)^i}{i} \binom{n-i}{i-1} y_1^{n+1-2i} z_1^i.$$

Then, we have

$$\begin{aligned} \frac{\partial f(y_1, z_1)}{\partial y_1} &= (n+1)y_1^n + (n+1) \sum_{1 \leq 2i \leq n} \frac{(-1)^i(n+1-2i)}{i} \binom{n-i}{i-1} y_1^{n-2i} z_1^i, \\ \frac{\partial f(y_1, z_1)}{\partial z_1} &= (n+1) \sum_{1 \leq 2i \leq n+1} (-1)^i \binom{n-i}{i-1} y_1^{n+1-2i} z_1^{i-1}. \end{aligned}$$

In the case that n is even:

The sequence $(\frac{\partial f(y_1, z_1)}{\partial y_1}, \frac{\partial f(y_1, z_1)}{\partial z_1})$ can be described as

$$((n+1)y_1^n + (-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} + p(y_1, z_1), -(n+1)y_1^{n-1} + q(y_1, z_1)),$$

where the polynomial degree $p(y_1, z_1)$ for y_1 satisfies

$$\deg_{y_1}(p(y_1, z_1)) < n-1$$

and the polynomial degree for z_1 satisfies

$$\deg_{z_1}(p(y_1, z_1)) < \frac{n}{2}$$

and the polynomial $q(y_1, z_1)$ satisfies that

$$\deg_{y_1}(q(y_1, z_1)) < n - 2$$

and

$$\deg_{z_1}(q(y_1, z_1)) < \frac{n}{2}.$$

It is easy to find that this sequence is a regular sequence. By Lemma 2.3 and Lemma 2.8, we have

$$\begin{aligned} \mathcal{C}(\bigcirc \rightarrow (y_1, z_1))_n &\simeq K \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}; \begin{pmatrix} -(n+1)y_1^n - (-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} - p(y_1, z_1) \\ (n+1)y_1^{n-1} - q(y_1, z_1) \end{pmatrix} \right)_{\mathbb{Q}[y_1, z_1]} \{1-n\}\{3-n\} \\ &\quad \left(\mathbb{Q}[y_1, z_1] \longrightarrow 0 \longrightarrow \mathbb{Q}[y_1, z_1] \right) \\ &\simeq K \left(\begin{pmatrix} 0 \\ 0 \end{pmatrix}; \begin{pmatrix} -(n+1)y_1^n - (-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} - p(y_1, z_1) \\ (n+1)y_1^{n-1} - q(y_1, z_1) \end{pmatrix} \right)_{\mathbb{Q}[y_1, z_1]} \{4-2n\}. \end{aligned}$$

We apply Theorem 2.10 to the polynomial $(n+1)y_1^{n-1} - q(y_1, z_1)$ for y_1 . Then, we obtain

$$\begin{aligned} (\mathbb{Q}[y_1, z_1] / \langle (n+1)y_1^{n-1} - q(y_1, z_1) \rangle) &\longrightarrow 0 \longrightarrow \mathbb{Q}[y_1, z_1] / \langle (n+1)y_1^{n-1} - q(y_1, z_1) \rangle \\ &\quad \boxtimes \\ K \left(0; -(-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} - y_1 q(y_1, z_1) - p(y_1, z_1) \right) &\longrightarrow \mathbb{Q}[y_1, z_1] / \langle (n+1)y_1^{n-1} - q(y_1, z_1) \rangle \\ &\quad \{4-2n\}. \end{aligned}$$

Since we have $\mathbb{Q}[y_1, z_1] / \langle (n+1)y_1^{n-1} - q(y_1, z_1) \rangle \simeq \mathbb{Q}[y_1] / \langle y_1^{n-1} \rangle [z_1]$, $\deg_{z_1}(y_1 q(y_1, z_1) + p(y_1, z_1)) < \frac{n}{2}$ and the sequence $((n+1)y_1^n + (-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} + p(y_1, z_1), -(n+1)y_1^{n-1} + q(y_1, z_1))$ is regular, we can once apply Theorem 2.10 to the polynomial $-(-1)^{\frac{n}{2}}(n+1)z_1^{\frac{n}{2}} - y_1 q(y_1, z_1) - p(y_1, z_1)$ for z_1 . Hence, we obtain

$$\mathcal{C}(\bigcirc \rightarrow (y_1, z_1))_n \simeq \left(J_{f(y_1, z_1)} \{4-2n\} \longrightarrow 0 \longrightarrow J_{f(y_1, z_1)} \{4-2n\} \right).$$

In the case that n is odd:

The sequence $(\frac{\partial f(y_1, z_1)}{\partial y_1}, \frac{\partial f(y_1, z_1)}{\partial z_1})$ can be described as

$$((n+1)y_1^n + p(y_1, z_1), -(n+1)y_1^{n-1} + (-1)^{\frac{n+1}{2}}(n+1)z_1^{\frac{n-1}{2}} + q(y_1, z_1)),$$

where the polynomial degree $p(y_1, z_1)$ for y_1 satisfies

$$\deg_{y_1}(p(y_1, z_1)) < n - 1$$

and the polynomial degree for z_1 satisfies

$$\deg_{z_1}(p(y_1, z_1)) < \frac{n+1}{2}$$

and the polynomial $q(y_1, z_1)$ satisfies that

$$\deg_{y_1}(q(y_1, z_1)) < n - 2$$

and

$$\deg_{z_1}(q(y_1, z_1)) < \frac{n-1}{2}.$$

By Lemma 2.3 and Lemma 2.8, we have

$$\mathcal{C}'\left(\bigcirc \begin{array}{c} \nearrow \\ \searrow \end{array} (y_1, z_1)\right)_n \simeq K \left(\left(\begin{array}{c} 0 \\ 0 \end{array} \right); \left(\begin{array}{c} -(n+1)y_1^n - p(y_1, z_1) \\ (n+1)y_1^{n-1} - (-1)^{\frac{n+1}{2}}(n+1)z_1^{\frac{n-1}{2}} - q(y_1, z_1) \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1]} \{1-n\}\{3-n\}.$$

Now, we have that $\deg_{y_1}(p(y_1, z_1)) < n-1$, $\deg_{z_1}(q(y_1, z_1) - (n+1)y_1^{n-1}) < \frac{n-1}{2}$ and the sequence $(\frac{\partial f(y_1, z_1)}{\partial y_1}, \frac{\partial f(y_1, z_1)}{\partial z_1})$ is regular. Thus, after we apply Theorem 2.10 to $-(n+1)y_1^n - p(y_1, z_1)$ for y_1 , we can once apply Theorem 2.10 to

$(n+1)y_1^{n-1} - (-1)^{\frac{n+1}{2}}(n+1)z_1^{\frac{n-1}{2}} - q(y_1, z_1)$ for z_1 . We also obtain

$$\mathcal{C}'\left(\bigcirc \begin{array}{c} \nearrow \\ \searrow \end{array} (y_1, z_1)\right)_n \simeq \left(J_{f(y_1, z_1)}\{4-2n\} \longrightarrow 0 \longrightarrow J_{f(y_1, z_1)}\{4-2n\} \right).$$

(4) By definition, we have

$$\mathcal{C}'\left(\begin{array}{c} (y_1, z_1) \\ \uparrow \\ x_3 \text{ --- } \bigcirc \text{ --- } x_4 \\ \uparrow \\ (y_2, z_2) \end{array}\right)_n = K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(x_3 + x_4, z_1)}{y_1 - x_3 - x_4} \\ \frac{f(x_3 + x_4, z_1) - f(x_3 + x_4, x_3 x_4)}{z_1 - x_3 x_4} \\ \frac{f(x_3 + x_4, x_3 x_4) - f(y_2, x_3 x_4)}{x_3 + x_4 - y_2} \\ \frac{f(y_2, x_3 x_4) - f(y_2, z_2)}{x_3 x_4 - z_2} \end{array} \right); \left(\begin{array}{c} y_1 - x_3 - x_4 \\ z_1 - x_3 x_4 \\ x_3 + x_4 - y_2 \\ x_3 x_4 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1, y_2, z_2, x_3, x_4]} \{-1\}.$$

Since the potential of this matrix factorization does not include the parameter x_3 and x_4 , we can apply Theorem 2.10 to the polynomial $y_1 - x_3 - x_4$ for x_3 . Thus, this matrix factorization is equivalent to the following matrix factorization

$$K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_1, -x_4^2 + y_1 x_4)}{z_1 + x_4^2 - y_1 x_4} \\ \frac{f(y_1, -x_4^2 + y_1 x_4) - f(y_2, -x_4^2 + y_1 x_4)}{y_1 - y_2} \\ \frac{f(y_2, -x_4^2 + y_1 x_4) - f(y_2, z_2)}{-x_4^2 + y_1 x_4 - z_2} \end{array} \right); \left(\begin{array}{c} z_1 + x_4^2 - y_1 x_4 \\ y_1 - y_2 \\ -x_4^2 + y_1 x_4 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1, y_2, z_2, x_3, x_4] / \langle y_1 - x_3 - x_4 \rangle} \{-1\}$$

in $\text{HMF}_{\mathbb{Q}[y_1, z_1, y_2, z_2, x_4], f(y_1, z_1) - f(y_2, z_2)}^{gr}$. Since the quotient ring $\mathbb{Q}[y_1, z_1, y_2, z_2, x_4] / \langle y_1 - x_3 - x_4 \rangle$ is isomorphic to

$$\mathbb{Q}[y_1, z_1, y_2, z_2, x_4],$$

we once apply Theorem 2.10 to the polynomial $z_1 + x_4^2 - y_1 x_4$ for x_4 . Then, this matrix factorization is equivalent to the following matrix factorization

$$K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_2, z_1)}{y_1 - y_2} \\ \frac{f(y_2, z_1) - f(y_2, z_2)}{z_1 - z_2} \end{array} \right); \left(\begin{array}{c} y_1 - y_2 \\ z_1 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1, y_2, z_2, x_4] / \langle z_1 + x_4^2 - y_1 x_4 \rangle} \{-1\}$$

in $\text{HMF}_{\mathbb{Q}[y_1, z_1, y_2, z_2], f(y_1, z_1) - f(y_2, z_2)}^{gr}$. The quotient ring $\mathbb{Q}[y_1, z_1, y_2, z_2, x_4] / \langle z_1 + x_4^2 - y_1 x_4 \rangle \{-1\}$ is isomorphic to

$$\mathbb{Q}[y_1, z_1, y_2, z_2] \{-1\} \oplus \mathbb{Q}[y_1, z_1, y_2, z_2] x_4 \{-1\}.$$

Furthermore, this ring is equivalent to

$$\mathbb{Q}[y_1, z_1, y_2, z_2] \{-1\} \oplus \mathbb{Q}[y_1, z_1, y_2, z_2] \{1\}$$

as a $\mathbb{Z}$ -graded $\mathbb{Q}[y_1, z_1, y_2, z_2]$ -module. Hence, we obtain the following matrix factorization

$$K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_2, z_1)}{y_1 - y_2} \\ \frac{f(y_2, z_1) - f(y_2, z_2)}{z_1 - z_2} \end{array} \right); \left(\begin{array}{c} y_1 - y_2 \\ z_1 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1, y_2, z_2]} \{-1\} \oplus K \left(\left(\begin{array}{c} \frac{f(y_1, z_1) - f(y_2, z_1)}{y_1 - y_2} \\ \frac{f(y_2, z_1) - f(y_2, z_2)}{z_1 - z_2} \end{array} \right); \left(\begin{array}{c} y_1 - y_2 \\ z_1 - z_2 \end{array} \right) \right)_{\mathbb{Q}[y_1, z_1, y_2, z_2]} \{1\}.$$

This is the right-hand side of the equivalent of Proposition (4).

(5) By definition, we have

$$\mathcal{C} \left(\begin{array}{ccc} x_1 & \longleftrightarrow & x_2 \\ & \downarrow (y_5, z_5) & \\ x_3 & \longleftrightarrow & x_4 \end{array} \right)_n = K \left(\left(\begin{array}{c} \frac{f(y_5, z_5) - f(x_3 + x_4, z_5)}{y_5 - x_3 - x_4} \\ \frac{f(x_3 + x_4, z_5) - f(x_3 + x_4, x_3 x_4)}{z_5 - x_3 x_4} \\ \frac{f(x_1 + x_2, x_1 x_2) - f(y_5, x_1 x_2)}{x_1 + x_2 - y_5} \\ \frac{f(y_5, x_1 x_2) - f(y_5, z_5)}{x_1 x_2 - z_5} \end{array} \right); \left(\begin{array}{c} y_5 - x_3 - x_4 \\ z_5 - x_3 x_4 \\ x_1 + x_2 - y_5 \\ x_1 x_2 - z_5 \end{array} \right) \right)_{\mathbb{Q}[x_1, x_2, x_3, x_4, y_5, z_5]} \{-1\}.$$

Since the potential of this matrix factorization does not include the parameter y_5 and z_5 , we apply Theorem 2.10 to the polynomials $y_5 - x_3 - x_4$ and $z_5 - x_3 x_4$. Then, it is easy to find that we obtain the right-hand side of the equivalent of Proposition (5). $\square$

A matrix factorization for the double loop $\bigcirc \bigcirc$ is defined as the above matrix factorization;

$$\mathcal{C}(\bigcirc \bigcirc)_n = (J_{f(y, z)}\{4 - 2n\} \longrightarrow 0 \longrightarrow J_{f(y, z)}\{4 - 2n\}).$$

Since the potential of the matrix factorization $\mathcal{C}(\bigcirc \bigcirc)_n$ is 0, we can take the homology group H of this matrix factorization.

Corollary 3.6. *The Euler characteristic of the homology $H(\mathcal{C}(\bigcirc \bigcirc)_n)$ equals the value of $\mathfrak{sl}_n$ quantum link invariant for the double loop;*

$$\bar{\chi}(H(\mathcal{C}(\bigcirc \bigcirc)_n)) = \frac{[n][n-1]}{[2]}.$$

Proof. By proof of the above Proposition 3.5 (3), we have

$$J_{f(y, z)} \simeq \begin{cases} \mathbb{Q}[y, z] / \langle y^{n-1}, z^{\frac{n}{2}} \rangle & n : \text{even} \\ \mathbb{Q}[y, z] / \langle y^n, z^{\frac{n-1}{2}} \rangle & n : \text{odd} \end{cases}$$

(i) $n : \text{even}$

Since $\deg_{\mathbb{Z}}(y_1) = 2$ and $\deg_{\mathbb{Z}}(z_1) = 4$, we have

$$\begin{aligned} \bar{\chi}(H(\mathcal{C}(\bigcirc \bigcirc)_n)) &= \bar{\chi}(J_{f(y, z)}\{4 - 2n\}) \\ &= \sum_{i=0}^{n-2} \sum_{j=0}^{\frac{n-2}{2}} q^{4-2n+2i+4j} \\ &= \sum_{i=0}^{n-2} q^{2-n+2i} \sum_{j=0}^{\frac{n-2}{2}} q^{2-n+4j} \\ &= (q^{2-n} + q^{4-n} + q^{6-n} + \cdots + q^{n-4} + q^{n-2})(q^{2-n} + q^{6-n} + q^{10-n} + \cdots + q^{n-6} + q^{n-2}) \\ &= \frac{[n][n-1]}{[2]}. \end{aligned}$$

(ii) n : odd

We have

$$\begin{aligned}
\overline{\chi}\left(H(\mathcal{C}(\bigcirc_n))\right) &= \overline{\chi}(J_{f(y,z)}\{4-2n\}) \\
&= \sum_{i=0}^{n-1} \sum_{j=0}^{\frac{n-3}{2}} q^{4-2n+2i+4j} \\
&= \sum_{i=0}^{n-1} q^{1-n+2i} \sum_{j=0}^{\frac{n-3}{2}} q^{3-n+4j} \\
&= (q^{1-n} + q^{3-n} + q^{5-n} + \cdots + q^{n-3} + q^{n-1})(q^{3-n} + q^{7-n} + q^{11-n} + \cdots + q^{n-7} + q^{n-3}) \\
&= \frac{[n][n-1]}{[2]}.
\end{aligned}$$

□

REFERENCES

- [1] D. Bar-Natan, Khovanov's homology for tangles and cobordisms, *Geom. Topol.* **9** (2005), 1443–1499 .
- [2] B. Gornik, Note on Khovanov link cohomology, [arXiv:math.QA/0402266](#).
- [3] S. Gukov, J. Walcher, Matrix Factorizations and Kauffman Homology, [arXiv:hep-th/0512298](#).
- [4] V. F. R. Jones, A polynomial invariant for knots via von Neumann algebras, *Bull. Amer. Math. Soc.* **12** (1985), 103–111.
- [5] L. Kauffman, State models and the Jones polynomial, *Topology*. **26** (1987), 395–407.
- [6] M. Khovanov, A categorification of the Jones polynomial, *Duke Math. J.* **101** (2000), no. 3, 359–426.
- [7] M. Khovanov, L. Rozansky, Matrix factorizations and link homology, [arXiv:math.QA/0401268](#).
- [8] M. Khovanov, L. Rozansky, Matrix factorizations and link homology II, [arXiv:math.QA/0505056](#).
- [9] M. Khovanov, L. Rozansky, Virtual crossings, convolutions and a categorification of the $SO(2N)$ Kauffman polynomial, [arXiv:math.QA/0701333](#)
- [10] G. Kuperberg, Spiders for rank 2 Lie algebras, *Comm. Math. Phys.* **180** (1996), no.1, 109–151.
- [11] H. Murakami, T. Ohtsuki, S. Yamada, Homfly polynomial via an invariant of colored plane graphs, *Enseign. Math.* (2) **44** (1998), no. 3-4, 325–360.
- [12] J. Rasmussen, Some differentials on Khovanov-Rozansky homology, [arXiv:math.GT/0607544](#).
- [13] Y. Yonezawa, in preparation.
- [14] B. Webster, Khovanov-Rozansky homology via a canopolis formalism, [arXiv:math.GT/0610650](#).
- [15] H. Wu, On the quantum filtration of the Khovanov-Rozansky cohomology, [arXiv:math.GT/0612406](#).

GRADUATE SCHOOL OF MATHEMATICS, NAGOYA UNIVERSITY, 464-8602 FUROCHO, CHIKUSAKU, NAGOYA, JAPAN
E-mail address: m03039e@math.nagoya-u.ac.jp